

IOWA STATE UNIVERSITY
Digital Press

Animal Industry Report

Animal Industry Report

AS 662

ASL R3120

2016

Does Heat Stress Alter the Pig's Response to Dietary Fat Source, as it Relates to Apparent or True Total Track Digestibility?

Trey A. Kellner

Iowa State University, tkellner@iastate.edu

Lance H. Baumgard

Iowa State University, baumgard@iastate.edu

Ken J. Prusa

Iowa State University, kprusa@iastate.edu

John F. Patience

Iowa State University, jfp@iastate.edu

Follow this and additional works at: https://lib.dr.iastate.edu/ans_air



Part of the [Agriculture Commons](#), [Food Science Commons](#), and the [Meat Science Commons](#)

Recommended Citation

Kellner, Trey A.; Baumgard, Lance H.; Prusa, Ken J.; and Patience, John F. (2016) "Does Heat Stress Alter the Pig's Response to Dietary Fat Source, as it Relates to Apparent or True Total Track Digestibility?," *Animal Industry Report*. AS 662, ASL R3120.

DOI: https://doi.org/10.31274/ans_air-180814-250

Available at: https://lib.dr.iastate.edu/ans_air/vol662/iss1/81

This Swine is brought to you for free and open access by the Animal Science Research Reports at Iowa State University Digital Repository. It has been accepted for inclusion in Animal Industry Report by an authorized editor of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Does Heat Stress Alter the Pig's Response to Dietary Fat Source, as it Relates to Apparent or True Total Tract Digestibility?

A.S. Leaflet R3120

Trey A. Kellner, Research Associate;
Lance H. Baumgard, Professor, Department of Animal
Science;

Ken J. Prusa, Professor, Department of Food Science and
Human Nutrition;

John F. Patience, Professor, Department of Animal Science

Summary and Implications

Heat stress affects a plethora of pork production variables, in part stemming from a reduction of feed intake. The experimental objective was to investigate the effect of heat stress on the pig's response to dietary fat in terms of growth performance and digestibility over a 35 d finishing period. A total of 96 barrows were randomly allotted to 1 of 9 treatments arranged as a 3×3 factorial with the main effects of environment [thermonetural (TN), pair-fed thermoneutral (PFTN), or heat stress (HS)] and diet [a corn-soybean meal based diet with 0% added fat (CNTR), or the CNTRL with 3% added tallow (3%TAL), or 3% added corn oil (3%CO)]. Pigs were individually housed to record intake. Fecal samples were collected on d 17 (~ 114 kg). No significant interactions between environment and diet were observed ($P > 0.100$). HS decreased ADFI (27.8%; $P < 0.001$), ADG (HS = 0.72, TN = 1.03, PFTN = 0.78 kg/d; $P < 0.001$), and G:F (HS = 0.290, TN = 0.301, PFTN = 0.319; $P = 0.006$). G:F but not ADG or ADFI tended to increase with added fat (CNTR = 0.292, 3%TAL = 0.303, 3%CO = 0.314 g/100 g; $P \leq 0.073$). Environment had no impact of TTTD of AEE ($P = 0.118$). In summary, HS decreased ADFI, ADG, G:F and ATTD of AEE, but had no significant impact on TTTD of AEE. Therefore, the pig's response to dietary fat source is not different in heat stress conditions as compared to thermoneutral conditions.

Introduction

Heat stress results in major losses to the pork industry through a plethora of production variables. Dietary fat is included in swine diets during seasonally warm conditions to minimize the heat of digestion and to maintain energy intake. The experimental objective was to investigate the effect of heat stress on the pig's response to dietary fat in terms of growth performance and digestibility of apparent

(ATTD) and true total tract digestibility (TTTD) of acid hydrolyzed ether extract (AEE) over a 35 d finishing period.

Materials and Methods

A total of 96 barrows (PIC 337 \times C22/29) with an initial BW of 100.4 ± 1.2 kg were randomly allotted to 1 of 9 treatments arranged as a 3×3 factorial, with the main effects of environment [TN (thermonetural: constant 24°C; ad libitum access to feed), PFTN (pair-fed thermoneutral: constant 24°C; limit-fed based on previous HS daily feed intake), or HS (heat stress: cyclical 28°C nighttime, 33°C-35°C daytime; ab libitum access to feed)] and diet [a corn-soybean meal based diet with 0% added fat (CNTR), 3% added tallow (3%TAL; iodine value = 41.8), or 3% added corn oil (3%CO; iodine value = 123.0)]. Pigs were individually housed to record intake. Titanium dioxide was included at 0.4% as an indigestible marker. Fecal samples were collected on d 17 (~ 114 kg). TTTD (%) of AEE was calculated by correcting ATTD of AEE for endogenous fat losses at 20 g of AEE/kg of dry matter intake. Data were analyzed using PROC MIXED with environment and dietary treatment as fixed effects, and replicate (2 replicates of 48 barrows) as a random effect.

Results and Discussion

No significant interactions between environment and diet were observed ($P > 0.100$; Table 1). Rectal temperature (HS = 39.0, TN = 38.1, PFTN = 38.2°C) increased due to HS ($P < 0.001$). HS decreased ADFI (27.8%; $P < 0.001$), ADG (HS = 0.72, TN = 1.03, PFTN = 0.78 kg/d; $P < 0.001$), and G:F (HS = 0.290, TN = 0.301, PFTN = 0.319; $P = 0.006$). G:F but not ADG or ADFI tended to increase with added fat (CNTR = 0.292, 3%TAL = 0.303, 3%CO = 0.314 g/100 g; $P \leq 0.073$). HS tended to have the lowest ATTD of AEE (HS = 59.0, TN = 60.2, PFTN = 61.4%, $P = 0.055$). Inclusion of dietary fat, and a source that was unsaturated increased ATTD of AEE (CNTR = 41.6, 3%TAL = 67.9, 3%CO = 71.2%, $P < 0.001$). TTTD of AEE of 3%CO-based diets was higher (99.3%) than that of CNTR (97.3%) and 3%TAL-based diets (96.3%; $P = 0.012$). Environment had no impact of TTTD of AEE ($P = 0.118$). In summary, HS decreased ADFI, ADG, G:F and ATTD of AEE, but had no significant impact on TTTD of AEE. Therefore, the pig's response to dietary fat source is not affected by HS. The inclusion of dietary fat cannot fully overcome the negative effects of HS.

Acknowledgements

We gratefully acknowledge the National Pork Board for funding this experiment, along with, Feed Energy Co., DSM Nutritional Products, and Anjinimoto Heartland for financial and other in-kind support or contributions.

Iowa State University Animal Industry Report 2016

Table 1. Effects of ad-libitum feed intake in thermal neutral conditions (TN)¹, pair-feeding in thermal neutral conditions (PFTN)¹, or heat stress (HS)² and additional inclusion of no dietary fat (CNTR), 3% tallow (TAL), or 3% corn oil (CO) on growth performance, rectal temperature³, respiration rate³, apparent total track digestibility (ATTD)⁴, and true total track digestibility (TTTD)⁴ of acid hydrolyzed ether extract (AEE)

Parameter	Treatments									P value ⁵				
	TN			PFTN			HS			SEM	T	E	DF	E × DF
	CNTR	TAL	CO	CNTR	TAL	CO	CNTR	TAL	CO					
ADG, kg	1.00 ^a	0.99 ^a	1.10 ^a	0.78 ^b	0.80 ^b	0.74 ^b	0.72 ^b	0.69 ^b	0.76 ^b	0.04	<0.001	<0.001	0.491	0.413
ADFI, kg	3.58 ^a	3.35 ^a	3.44 ^a	2.54 ^b	2.42 ^b	2.52 ^b	2.54 ^b	2.41 ^b	2.51 ^b	0.13	<0.001	<0.001	0.124	0.978
G:F, kg	0.287 ^{bc}	0.295 ^{bc}	0.320 ^{ab}	0.307 ^{abc}	0.332 ^a	0.317 ^{ab}	0.282 ^c	0.284 ^c	0.304 ^{abc}	0.016	0.011	0.006	0.073	0.500
RR ⁶ , bpm	37.1 ^b	36.8 ^b	35.0 ^b	34.8 ^b	34.5 ^b	33.8 ^b	79.3 ^a	77.4 ^a	78.1 ^a	2.1	<0.001	<0.001	0.692	0.904
RT ⁷ , °C	38.1 ^b	38.1 ^b	38.2 ^b	38.2 ^b	38.2 ^b	38.2 ^b	39.0 ^a	39.0 ^a	38.9 ^a	0.1	<0.001	<0.001	0.653	0.191
ATTD, %	41.3 ^e	67.9 ^{bcd}	71.5 ^{ab}	42.9 ^{de}	68.4 ^{bcd}	72.8 ^a	40.4 ^e	67.3 ^{cd}	69.4 ^{bc}	1.3	<0.001	0.054	<0.001	0.886
TTTD ⁸ , %	97.1	96.4	100.1	98.8	96.8	99.9	96.2	95.7	98.0	0.7	0.082	0.118	0.012	0.932

^{a-c}Within a row, least squares means lacking a common superscript differ, $P < 0.05$.

¹Constant thermal neutral environment of ~24.0°C.

²Diurnal heat stress environment of ~33.0°C between 0800 h to 2000 h and ~28.0°C 2000 h to 0800 h from d 0 to d 7, ~33.5°C between 0800 h to 2000 h and ~28.0°C 2000 h to 0800 h for d 7 to d 14, ~34.0°C between 0800 h to 2000 h and ~28.0°C 2000 h to 0800 h for d 14 to d 21, ~34.5°C between 0800 h to 2000 h and ~28.0°C 2000 h to 0800 h for d 21 to d 28, and ~35.0°C between 0800 h to 2000 h and ~28.0°C 2000 h to 0800 h for d 28 to d 35.

³Measured daily at 1100 h.

⁴Measured on d 17.

⁵Probability values for main effects of treatment (T), environment (E), and dietary fat (DF), as well as the environment × dietary fat interaction (E × DF).

⁶Respiration rate (breaths per minute).

⁷Rectal temperature (°C).

⁸TTTD (%) of AEE was calculated by correcting ATTD of AEE for endogenous fat losses at 20 g of AEE/kg of dry matter intake.